

JEREMY S. LEWIS, PT, PhD, FCSP^{1,3} • CHAD E. COOK, PT, MBA, PhD, FAPTA^{4,5}
TAMMY C. HOFFMANN, PhD⁶ • PETER O'SULLIVAN, PT, PhD^{7,8}

The Elephant in the Room: Too Much Medicine in Musculoskeletal Practice

J Orthop Sports Phys Ther 2020;50(1):1-4. doi:10.2519/jospt.2020.0601

Advances in assessment and management of musculoskeletal conditions (eg, fracture management) have improved care for many people. We contend that there have been other, less beneficial developments in the provision of care for people with musculoskeletal pain conditions—one is the worrying tendency to provide too much medicine.

There are overlaps and confusion regarding the usage and definitions of terms.⁸ In this Viewpoint, we will use the term “too much medicine” as an umbrella term that includes overdiagnosis, misdiagnosis, false positives, diagnostic overmedicalization, and overdetection. Too much medicine has led to overtreatment, overutilization, interventional overmedicalization, and low-value care.¹²

Many musculoskeletal conditions require a level of investigation and intervention. Too much medicine occurs when the provision of either (or both) is unjustifiably excessive, for example, referring an individual experiencing non-specific low back pain with no red flags for magnetic resonance imaging. Another concern in musculoskeletal health care is

medicalizing normality—when a normal human function or condition is labeled as abnormal.¹⁰

In this Viewpoint, we argue that too much medicine and medicalizing normality in contemporary musculoskeletal practice have become the “elephant in the room.” Medicalizing normality creates health concerns where none exist. Too much medicine involves provision of care where benefits do not outweigh harms, and wastes precious health care resources. We (1) list 2 common examples of too much medicine, and 2 examples of medicalizing normality, relevant to physical therapy practice; (2) outline the drivers of too much medicine and medicalizing normality; and (3) make suggestions for change.

Two Examples of Too Much Medicine Relevant to Physical Therapy Practice

Nonsurgical Interventions for Pain Musculoskeletal pain management costs continue to rise. Individuals may have been misinformed that myriad nonsurgical health care options, including acupuncture, manual therapy, myofascial trigger point therapy, injections, pharmacology, among others, will, in isolation, “fix” the problem. Use of opioids has been at the forefront of the drive to eradicate pain. Worldwide, use of prescription opioid analgesics more than doubled between 2001 and 2013, leading to an opioid epidemic in many countries.¹⁴ In the United States in 2017, health care providers (principally general practitioners) prescribed opioid pain medication 191 million times (59 prescriptions per 100 people). Twenty-nine million people were taking nonsteroidal anti-inflammatory medications, accompanied by 100 000 hospitalizations and 17 000 related deaths.^{4,19}

Orthopaedic Surgery Many surgical procedures perform no better than skin in-

¹School of Health and Social Work, University of Hertfordshire, Hatfield, United Kingdom. ²Therapy Department, Central London Community Healthcare National Health Services Trust, London, United Kingdom. ³Department of Physical Therapy and Rehabilitation Science, College of Health Sciences, Qatar University, Doha, Qatar. ⁴Doctor of Physical Therapy Division, Duke University School of Medicine, Durham, NC. ⁵Physical Therapy Division and Duke Clinical Research Institute, Duke University, Durham, NC. ⁶Institute for Evidence-Based Healthcare, Faculty of Health Sciences and Medicine, Bond University, Gold Coast, Australia. ⁷School of Physiotherapy and Exercise Science, Health Sciences Division, Curtin University, Bentley, Australia. ⁸Body Logic Physiotherapy, Shenton Park, Australia. No funding was received to support this Viewpoint. The authors certify that they have no affiliations with or financial involvement in any organization or entity with a direct financial interest in the subject matter or materials discussed in the article. Address correspondence to Dr Jeremy Lewis, School of Health and Social Work, Wright Building, College Lane Campus, University of Hertfordshire, Hatfield AL10 9AB Hertfordshire, UK. E-mail: jeremy.lewis@LondonShoulderClinic.com • Copyright ©2020 *Journal of Orthopaedic & Sports Physical Therapy*[®]

cisions and arthroscopy without the “fix” (sham surgery), especially when the patient’s main complaint is pain.⁷ Examples include repairs for nontraumatic medical meniscal tears, arthroscopic debridement for knee osteoarthritis,¹³ type II superior labral tear from anterior to posterior lesions,²⁰ biceps tenodesis for long-head-of-biceps pathology,²⁰ and acromioplasty for subacromial impingement.¹⁶

Prioritizing expensive surgical procedures when cheaper, equally effective alternatives exist is concerning. Non-surgical management, principally in the form of graduated activity and exercise, is consistently as effective as surgery for shoulder pain, knee pain, and the majority of grade I to III ankle sprains. Surgery may be a reasonable treatment option, but it is associated with increased clinical risk and increased costs for health systems and patients, often without providing increased clinical benefits. For many musculoskeletal conditions, too much medicine can be avoided if appropriate condition-specific education, lifestyle advice, and evidence-based non-surgical management are prioritized.

Two Examples of Medicalizing Normality Relevant to Physical Therapy Practice

Musculoskeletal aches and pains are common. Up to 70% of people experience shoulder pain and 90% experience low back pain at some stage in their lives. One might argue that these common musculoskeletal conditions could be considered unpleasant yet “normal” occurrences. In this section, we summarize 2 examples of the mislabeling of normal and age-related variations in posture and structure as “pathological” and/or the basis for presenting symptoms.

Postural “Abnormalities” Ankylosing spondylitis and severe kyphosis and scoliosis may be associated with symptoms. However, for the majority of musculoskeletal presentations, most posture “abnormalities” are likely to be variations of normal and do not differentiate between people with and without pain.¹¹ Observing a person’s static posture based on the

plumb-line assessment of cervical, thoracic, and shoulder posture, then advising that the symptoms are due to subtle variations in postural alignment, is medicalizing normality.

“Abnormalities” Detected by Imaging There has been an increase in the identification of “abnormalities” in magnetic resonance imaging and ultrasound as the explanation for presenting symptoms. However, this practice has medicalized normality on an unprecedented scale.^{1,2} Examples include lumbar disc protrusions, disc bulges, facet joint degeneration, and spondylolisthesis in people without low back pain; labral abnormalities and rotator cuff tendon pathology in baseball pitchers without shoulder pain; osteophytes, cartilage damage, bone marrow lesions, and synovitis in people without knee pain; and labral tears in young people without hip pain. These findings suggest that many changes labeled as “abnormalities” are normal and may not be associated with pain or symptoms. Many interventions may be performed on people who have normal age-related changes, and most probably on tissues that are not the cause of the symptoms.

Drivers of Too Much Medicine

There are many drivers of too much medicine, including the belief of clinicians and patients that more health care (in the form of imaging and investigations, prescribing medicine, injections, multiple passive interventions and electrotherapy modalities, and surgery) is better than

prioritizing condition-specific and lifestyle advice; that “doing something” is better than “waiting and watching”⁶; that the origin of pain can always be identified with clinical tests and imaging; that once identified, pain can be “fixed”; that symptoms are caused by “abnormalities” in static posture and structure; and that not addressing “abnormalities” risks further tissue damage or exacerbation of the condition.³

When more expensive interventions are recommended that offer equivalent or worse outcomes than lower-cost alternatives, profit and remuneration become drivers of sectors of the health care industry, insurers, pharmaceutical companies, and some clinicians. For sections of the media, drivers include sensationalism and revenue.⁹ Politicians may not wish to disenfranchise voters by appearing to reduce or withdraw health care alternatives considered fundamental by the electorate or advocated by lobby groups.^{15,18}

Suggestions for Change

Reducing the sequelae of too much medicine will require continuous effort from all stakeholders.¹⁷ We must all consider sustainability and acknowledge that health care resources are finite. In this section, we outline suggestions for how patients, policy makers, clinicians, educators, the health care industry, and the media can drive change. We have included a recommended reading list (**APPENDIX**, available at www.jospt.org) and resources (**TABLE**) to help reduce the impact of too

TABLE

RESOURCES TO BETTER UNDERSTAND AND HELP REDUCE TOO MUCH MEDICINE AND MEDICALIZING NORMALITY

- Provide a card or leaflet with possible questions or discussion points for patients: <https://www.nhs.uk/using-the-nhs/nhs-services/gps/what-to-ask-your-doctor/>
- Discuss the value and importance of shared decision making in musculoskeletal practice⁶
- Too much medicine: <https://www.bmj.com/too-much-medicine> and https://www.youtube.com/watch?v=FDfclD_BsA
- Better medicine: <https://www.bmj.com/bettermedicine>
- Improving critical thinking about health care: <https://www.healthnewsreview.org>
- Online guides for reporting medical research: <https://www.smh.com.au/national/our-guidelines-for-reporting-medical-research-20190603-p51tw2.html>
- Discuss musculoskeletal management in social media forums⁵

much medicine and medicalizing normality in musculoskeletal practice.

What Can Patients Do?

- Ask questions relating to the different management options for your condition, focusing on the anticipated benefits, time scales, and harms
- Ask what you can do to help manage your condition
- Ask if “wait and watch” is an appropriate option
- When fully informed of the benefits, harms, and costs of the management options, contribute to codesigning the provision of health care at local and national levels
- Share experiences and journeys through the health care system to promote improvements in delivery of care

What Can Policy Makers Do?

- Withdraw the financial incentive to offer unnecessary assessment and intervention
- Defund low-value care (eg, subacromial decompression surgery¹⁶)
- Prioritize funding high-value care (eg, exercise programs for rotator cuff-related shoulder pain⁶)

What Can Clinicians Do?

- Ensure patients are aware of and understand all reasonable diagnosis and management options, and the harms, benefits, and expected outcomes of each
- Avoid emotive language and outdated explanations when explaining symptoms and making recommendations for management
- Establish what matters most to the patient and discuss this as part of decision making
- Understand the natural course of the condition
- Know the investigations that should and should not be considered, and age-related norms for investigation findings
- Discuss the impact an intervention may have on the individual (eg, the number of sessions a patient may need to attend for an exercise class and how much self-directed exercise he or she would need to complete, any limita-

tions to activity following an injection or surgery, and for how long)

What Can Educators Do?

- Ensure that curricula are contemporary and reflect current evidence
- Teach critical appraisal skills so clinicians can effectively and efficiently incorporate new evidence into practice
- Teach shared decision-making skills

What Can the Health Care Industry Do?

- Use common language and explanations for patients, based on an unbiased assessment of research
- Promote interprofessional practice

What Can the Media Do?

- Recognize the harm and the distress that can occur when health care issues are sensationalized or misrepresented

SUMMARY

TOO MUCH MEDICINE BURDENS health care systems and deprives societies of resources. Overcoming too much medicine requires stakeholders (patients, clinicians, educators, health care funders, media, policy makers, industry, insurers, politicians, etc) to appropriately prioritize low-risk, cost-effective care over higher-risk and more expensive care of equal clinical effectiveness. Professional bodies, government agencies, clinicians, and patients must collaborate to discuss and synthesize the available evidence, share decisions, and translate knowledge. We do not underestimate the challenge of the task. Reducing the harm of medicalizing normality and avoiding the temptation to provide too much medicine in current musculoskeletal practice depend on all of us. ●

ACKNOWLEDGMENTS: *The authors wish to sincerely thank Dr Clare Ardern for her expert help and guidance in the preparation of this manuscript.*

REFERENCES

1. Barreto RPG, Braman JP, Ludewig PM, Ribeiro LP, Camargo PR. Bilateral magnetic resonance imaging findings in individuals with unilateral shoulder

1. *J Shoulder Elbow Surg.* 2019;28:1699-1706. <https://doi.org/10.1016/j.jse.2019.04.001>
2. Brinjikji W, Luetmer PH, Comstock B, et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol.* 2015;36:811-816. <https://doi.org/10.3174/ajnr.A4173>
3. Bunzli S, Smith A, Schütze R, O'Sullivan P. Beliefs underlying pain-related fear and how they evolve: a qualitative investigation in people with chronic back pain and high pain-related fear. *BMJ Open.* 2015;5:e008847. <https://doi.org/10.1136/bmjopen-2015-008847>
4. Centers for Disease Control and Prevention. U.S. opioid prescribing rate maps. Available at: <https://www.cdc.gov/drugoverdose/maps/rxrate-maps.html>. Accessed October 21, 2019.
5. Cook CE, O'Connell NE, Hall T, et al. Benefits and threats to using social media for presenting and implementing evidence. *J Orthop Sports Phys Ther.* 2018;48:3-7. <https://doi.org/10.2519/jospt.2018.0601>
6. Hoffmann TC, Lewis J, Maher CG. Shared decision making should be an integral part of physiotherapy practice. *Physiotherapy.* In press. <https://doi.org/10.1016/j.physio.2019.08.012>
7. Jonas WB, Crawford C, Colloca L, et al. To what extent are surgery and invasive procedures effective beyond a placebo response? A systematic review with meta-analysis of randomised, sham controlled trials. *BMJ Open.* 2015;5:e009655. <https://doi.org/10.1136/bmjopen-2015-009655>
8. Kaczmarek E. How to distinguish medicalization from over-medicalization? *Med Health Care Philos.* 2019;22:119-128. <https://doi.org/10.1007/s11019-018-9850-1>
9. Kilgo DK, Harlow S, García-Perdomo V, Salaverría R. A new sensation? An international exploration of sensationalism and social media recommendations in online news publications. *Journalism.* 2018;19:1497-1516. <https://doi.org/10.1177/1464884916683549>
10. Lewis J. The medicalisation of normality in musculoskeletal practice [abstract]. *J Sci Med Sport.* 2017;20 suppl 3:37. <https://doi.org/10.1016/j.jsams.2017.09.266>
11. Lewis JS, Green A, Wright C. Subacromial impingement syndrome: the role of posture and muscle imbalance. *J Shoulder Elbow Surg.* 2005;14:385-392. <https://doi.org/10.1016/j.jse.2004.08.007>
12. Mafi JN, Parchman M. Low-value care: an intractable global problem with no quick fix. *BMJ Qual Saf.* 2018;27:333-336. <https://doi.org/10.1136/bmjqs-2017-007477>
13. Moseley JB, O'Malley K, Petersen NJ, et al. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med.* 2002;347:81-88. <https://doi.org/10.1056/NEJMoa013259>
14. Neuman MD, Bateman BT, Wunsch H. Inappropriate opioid prescription after surgery. *Lancet.* 2019;393:1547-1557. [https://doi.org/10.1016/S0140-6736\(19\)30428-3](https://doi.org/10.1016/S0140-6736(19)30428-3)
15. Newton-Small J. The first 2020 Democratic primary debate will almost certainly skip a key

[VIEWPOINT]

healthcare issue. *Time*. June 26, 2019. Available at: <https://time.com/5614083/2020-democratic-debate-healthcare/>

16. Paavola M, Malmivaara A, Taimela S, et al. Subacromial decompression versus diagnostic arthroscopy for shoulder impingement: randomised, placebo surgery controlled clinical trial. *BMJ*. 2018;362:k2860. <https://doi.org/10.1136/bmj.k2860>
17. Pathirana T, Clark J, Moynihan R. Mapping the drivers of overdiagnosis to potential solutions. *BMJ*. 2017;358:j3879. <https://doi.org/10.1136/bmj.j3879>

bmj.j3879

18. Pearl R. Healthcare is the no. 1 issue for voters; a new poll reveals which healthcare issue matters most. *Forbes*. August 13, 2018. Available at: <https://www.forbes.com/sites/robertpearl/2018/08/13/midterms/#483775bc3667>
19. Rane MA, Foster JG, Wood SK, Hebert PR, Hennekens CH. Benefits and risks of nonsteroidal anti-inflammatory drugs: methodologic limitations lead to clinical uncertainties. *Ther Innov Regul Sci*. 2019;53:502-505. <https://doi.org/10.1177/2168479018794159>

20. Schrøder CP, Skare Ø, Reikerås O, Mowinckel P, Brox JI. Sham surgery versus labral repair or biceps tenodesis for type II SLAP lesions of the shoulder: a three-armed randomised clinical trial. *Br J Sports Med*. 2017;51:1759-1766. <https://doi.org/10.1136/bjsports-2016-097098>



MORE INFORMATION
WWW.JOSPT.ORG

PUBLISH Your Manuscript in a Journal With International Reach

JOSPT offers authors of accepted papers an **international audience**. The *Journal* is currently distributed to the members of APTA's Orthopaedic and Sports Physical Therapy Sections and **32 orthopaedics, manual therapy, and sports groups in 24 countries** who provide online access either as a member benefit or at a discount. As a result, the *Journal* is now distributed monthly to more than **37,000 individuals around the world** who specialize in musculoskeletal and sports-related rehabilitation, health, and wellness. In addition, *JOSPT* reaches students and faculty, physical therapists and physicians at more than **1,250 institutions in 60 countries**. Please review our Information for and Instructions to Authors at www.jospt.org in the **Info Center** for **Authors** and submit your manuscript for peer review at <http://mc.manuscriptcentral.com/jospt>.

APPENDIX

RECOMMENDED READING LIST

- Blendon RJ, Benson JM, McMurtry CL. Health care in the 2018 election. *N Engl J Med*. 2018;379:e32. <https://doi.org/10.1056/NEJMsri813425>
- Brodersen J, Kramer BS, Macdonald H, Schwartz LM, Woloshin S. Focusing on overdiagnosis as a driver of too much medicine. *BMJ*. 2018;362:k3494. <https://doi.org/10.1136/bmj.k3494>
- Brodersen J, Schwartz LM, Heneghan C, O'Sullivan JW, Aronson JK, Woloshin S. Overdiagnosis: what it is and what it isn't. *BMJ Evid Based Med*. 2018;23:1-3. <https://doi.org/10.1136/ebmed-2017-110886>
- Buchbinder R, van Tulder M, Öberg B, et al. Low back pain: a call for action. *Lancet*. 2018;391:2384-2388. [https://doi.org/10.1016/S0140-6736\(18\)30488-4](https://doi.org/10.1016/S0140-6736(18)30488-4)
- Carr AJ, Cooper CD, Campbell MK, et al. Clinical effectiveness and cost-effectiveness of open and arthroscopic rotator cuff repair [the UK Rotator Cuff Surgery (UKUFF) randomised trial]. *Health Technol Assess*. 2015;19:1-218. <https://doi.org/10.3310/hta19800>
- Carter SM, Rogers W, Heath I, Degeling C, Doust J, Barratt A. The challenge of overdiagnosis begins with its definition. *BMJ*. 2015;350:h869. <https://doi.org/10.1136/bmj.h869>
- Culvenor AG, Øiestad BE, Hart HF, Stefanik JJ, Guermazi A, Crossley KM. Prevalence of knee osteoarthritis features on magnetic resonance imaging in asymptomatic uninjured adults: a systematic review and meta-analysis. *Br J Sports Med*. 2019;53:1268-1278. <https://doi.org/10.1136/bjsports-2018-099257>
- Darlow B, Forster BB, O'Sullivan K, O'Sullivan P. It is time to stop causing harm with inappropriate imaging for low back pain. *Br J Sports Med*. 2017;51:414-415. <https://doi.org/10.1136/bjsports-2016-096741>
- GBD 2017 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1859-1922. [https://doi.org/10.1016/S0140-6736\(18\)32335-3](https://doi.org/10.1016/S0140-6736(18)32335-3)
- GBD 2017 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1789-1858. [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7)
- Guermazi A, Niu J, Hayashi D, et al. Prevalence of abnormalities in knees detected by MRI in adults without knee osteoarthritis: population based observational study (Framingham Osteoarthritis Study). *BMJ*. 2012;345:e5339. <https://doi.org/10.1136/bmj.e5339>
- Handoll H, Brealey S, Rangan A, et al. The ProFHER (PROximal Fracture of the Humerus: Evaluation by Randomisation) trial - a pragmatic multicentre randomised controlled trial evaluating the clinical effectiveness and cost-effectiveness of surgical compared with non-surgical treatment for proximal fracture of the humerus in adults. *Health Technol Assess*. 2015;19:1-280. <https://doi.org/10.3310/hta19240>
- Harradine P, Gates L, Bowen C. If it doesn't work, why do we still do it? The continuing use of subtalar joint neutral theory in the face of overpowering critical research. *J Orthop Sports Phys Ther*. 2018;48:130-132. <https://doi.org/10.2519/jospt.2018.0604>
- Hoffmann TC, Del Mar C. Clinicians' expectations of the benefits and harms of treatments, screening, and tests: a systematic review. *JAMA Intern Med*. 2017;177:407-419. <https://doi.org/10.1001/jamainternmed.2016.8254>
- Hoy D, Brooks P, Blyth F, Buchbinder R. The epidemiology of low back pain. *Best Pract Res Clin Rheumatol*. 2010;24:769-781. <https://doi.org/10.1016/j.berh.2010.10.002>
- Jonas WB, Crawford C, Colloca L, et al. To what extent are surgery and invasive procedures effective beyond a placebo response? A systematic review with meta-analysis of randomised, sham controlled trials. *BMJ Open*. 2015;5:e009655. <https://doi.org/10.1136/bmjopen-2015-009655>
- Kado DM, Huang MH, Karlamangla AS, Barrett-Connor E, Greendale GA. Hyperkyphotic posture predicts mortality in older community-dwelling men and women: a prospective study. *J Am Geriatr Soc*. 2004;52:1662-1667. <https://doi.org/10.1111/j.1532-5415.2004.52458.x>
- Lee AJ, Armour P, Thind D, Coates MH, Kang AC. The prevalence of acetabular labral tears and associated pathology in a young asymptomatic population. *Bone Joint J*. 2015;97-B:623-627. <https://doi.org/10.1302/0301-620X.97B5.35166>
- Lewis J. The end of an era? *J Orthop Sports Phys Ther*. 2018;48:127-129. <https://doi.org/10.2519/jospt.2018.0102>
- Lewis J, O'Sullivan P. Is it time to reframe how we care for people with non-traumatic musculoskeletal pain? *Br J Sports Med*. 2018;52:1543-1544. <https://doi.org/10.1136/bjsports-2018-099198>
- Luime JJ, Koes BW, Hendriksen IJ, et al. Prevalence and incidence of shoulder pain in the general population; a systematic review. *Scand J Rheumatol*. 2004;33:73-81. <https://doi.org/10.1080/03009740310004667>
- Miniaci A, Mascia AT, Salonen DC, Becker EJ. Magnetic resonance imaging of the shoulder in asymptomatic professional baseball pitchers. *Am J Sports Med*. 2002;30:66-73. <https://doi.org/10.1177/03635465020300012501>
- Moynihan R, Doust J, Henry D. Preventing overdiagnosis: how to stop harming the healthy. *BMJ*. 2012;344:e3502. <https://doi.org/10.1136/bmj.e3502>
- Moynihan R, Heath I, Henry D. Selling sickness: the pharmaceutical industry and disease mongering. *BMJ*. 2002;324:886-891. <https://doi.org/10.1136/bmj.324.7342.886>
- Pathirana T, Clark J, Moynihan R. Mapping the drivers of overdiagnosis to potential solutions. *BMJ*. 2017;358:j3879. <https://doi.org/10.1136/bmj.j3879>
- Petersen W, Rembitzki IV, Koppenburg AG, et al. Treatment of acute ankle ligament injuries: a systematic review. *Arch Orthop Trauma Surg*. 2013;133:1129-1141. <https://doi.org/10.1007/s00402-013-1742-5>
- Ransohoff DF, Ransohoff RM. Sensationalism in the media: when scientists and journalists may be complicit collaborators. *Eff Clin Pract*. 2001;4:185-188.
- Sihvonen R, Paavola M, Malmivaara A, et al. Arthroscopic partial meniscectomy versus placebo surgery for a degenerative meniscus tear: a 2-year follow-up of the randomised controlled trial. *Ann Rheum Dis*. 2018;77:188-195. <https://doi.org/10.1136/annrheumdis-2017-211172>
- Sokol DK. Update on the UK law on consent. *BMJ*. 2015;350:h1481. <https://doi.org/10.1136/bmj.h1481>
- van Dijk W, Faber MJ, Tanke MA, Jeurissen PP, Westert GP. Medicalisation and overdiagnosis: what society does to medicine. *Int J Health Policy Manag*. 2016;5:619-622. <https://doi.org/10.15171/ijhpm.2016.121>
- Vos T, Flaxman AD, Naghavi M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380:2163-2196. [https://doi.org/10.1016/S0140-6736\(12\)61729-2](https://doi.org/10.1016/S0140-6736(12)61729-2)